

Testing Electron Boost Invariance with $2S$ - $1S$ Hydrogen Spectroscopy

Brett Altschul¹

*Department of Physics and Astronomy
University of South Carolina
Columbia, SC 29208*

Abstract

There are few good direct laboratory tests of boost invariance for electrons, because the experiments required often involve repeated precision measurements performed at different times of year. However, existing measurements and remeasurements of the $2S$ - $1S$ two-photon transition frequency in ^1H —which were done to search for a time variation in the fine structure constant—also constitute a measurement of the boost symmetry violation parameter $0.83c_{(TX)} + 0.51c_{(TY)} + 0.22c_{(TZ)} = (4 \pm 8) \times 10^{-11}$. This is an eight order of magnitude improvement over preexisting laboratory bounds, and with only one additional measurements, this system could yield a second comparable constraint.

¹baltschu@physics.sc.edu

There is currently a great deal of interest in the possibility that Lorentz and *CPT* symmetries may not be exact in nature [1]. A diverse spectrum of precision experiments have placed bounds on many different forms of Lorentz violation. These various forms are described by the parameters of the standard model extension (SME), an effective quantum field theory [2]. Bounds on SME parameters are summarized in [3].

Atomic spectroscopy is a powerful tool in precision physics. Many of the most precise tests of Lorentz symmetry have used atomic clocks [4, 5, 6, 7, 8, 9]. These experiments mostly involve measurements of nuclear spin transitions, and less attention has been paid to optical transitions, which are typically sensitive to different sets SME coefficients. Optical transition frequencies are generally more sensitive to Lorentz violation in the electron sector and to forms of Lorentz violation that are independent of particle spin.

We shall show how existing optical spectroscopy measurements can be used to place new bounds on Lorentz violation. The measurements involved were of the $2S$ - $1S$ two-photon transition frequency in hydrogen [10, 11]. This has historically been one of the best known optical transition frequencies, and it provides useful information about many different physical phenomena. Lorentz violation in ^1H was previously discussed in [12]. However, since the focus there was primarily on frequency shifts that were not suppressed by any power of the fine structure constant α , the class of experiments described here was not discussed.

The electron Lagrange density relevant for the ^1H experiments is

$$\mathcal{L} = \bar{\psi}[(\gamma^\mu + c^{\nu\mu}\gamma_\nu)i\partial_\mu - m]\psi, \quad (1)$$

where $c^{\nu\mu}$ is a traceless background tensor describing the Lorentz violation. The electron sector of the SME contains other tensors, which parameterize different types of Lorentz and *CPT* violation. However, most of these depend on spin, and the ones that could affect the energy difference between two atomic S states are already strongly constrained by torsion pendulum experiments with spin-polarized samples [13].

The c_{00} term in $\mathcal{L}$ enters as a rescaling of the time derivative term in the action. Under quantization, $i\partial_0$ becomes the electron energy E , and c_{00} simply rescales all electronic energy eigenvalues. Including the effects of the all the $c^{\nu\mu}$ coefficients, the nonrelativistic Hamiltonian for the electron is [14]

$$H = \frac{p_j p_k}{2m} [\delta_{jk} - c_{00}\delta_{jk} - c_{(jk)}] - c_{(0j)}p_j + V. \quad (2)$$

The coefficients $c_{(\nu\mu)}$ are the symmetrized combinations $c_{\nu\mu} + c_{\mu\nu}$.

Bounds on SME coefficients are generally given in sun-centered celestial equatorial coordinates [15]. The sun-centered reference frame is approximately inertial on all relevant time scales, and it provides a convenient way to parameterize measurements of boost symmetry violation that make use of the Earth's orbital motion (such as the measurements discussed here). The Cartesian coordinates used in this frame are (X, Y, Z, T) . The spatial origin lies at the center of the sun. The Z -axis points along the direction of the Earth's

rotation, and the X -axis points toward the vernal equinox point on the celestial sphere. (This means Earth lies on the negative X -axis at the time of this equinox.) The Y -direction is chosen according to the right hand rule, and the origin of time ($T = 0$) is taken to be the vernal equinox in the year 2000.

There are no strong laboratory bounds on the $c_{(TJ)}$. Constraints based on Doppler effect measurements are only at the 10^{-2} level [16]; and even these weak bounds are only order of magnitude inferences based on earlier experimental data. The experiment used spectroscopy of ${}^7\text{Li}^+$ ions moving at a speed $v \approx 0.064$ to test the accuracy of the relativistic Doppler shift formula. The experiment confirmed the conventional prediction with $\sim 2 \times 10^{-9}$ precision [17], providing a weak sensitivity to the electron $c_{(TJ)}$ coefficients.

There are better astrophysical bounds [18], but they have a number of fairly undesirable features. These bounds are derived from observations of extremely high energy astrophysical phenomena. In order to translate the observational data into constraints on the $c_{(TJ)}$ and other coefficients, we must understand how the energetic phenomena really work. For example, the TeV γ -rays from many sources are probably produced by the upscattering of low-energy photons by extremely energetic electrons (inverse Compton scattering), and the photon spectrum can tell us a great deal about the behavior of the highly boosted electrons. However, there is an alternative hypothesis for the origins of the TeV photons—that they are produced in π^0 decay; if that is the case, then observations of the photons may really tell us very little about electron Lorentz violation.

Yet there is one type of astrophysical bound that is immune to this problem. However TeV γ -rays are produced, they could potentially decay ($\gamma \rightarrow e^+ + e^-$) if there is Lorentz violation in the electron sector. The observed absence of this process allows us to place some stringent constraints on the $c_{(TJ)}$ and c_{TT} coefficients [19]. However, these constraints are all one-sided, and it is impossible to disentangle the $c_{(TJ)}$ bounds from those on c_{TT} . Consequently, these measurements cannot exclude any specific values of $c_{(TJ)}$. This makes it extremely important to have separate, two-sided bounds on the $c_{(TJ)}$.

c_{00} acts to shift all electron energies, and its effects may be seen using any electron transition. Moreover, if one measures an observable that is even under C , T , and any reflection (the parity operator P may be decomposed into three separate reflections $P = R_1 R_2 R_3$), c_{00} and c_{jk} with $j = k$ are the only electron coefficient that can contribute at first order. All other forms of electron-sector Lorentz violation are odd under at least one of C , T , or R_j . The anisotropic c_{JK} coefficients can be constrained by looking at Michelson-Morley experiments with material-filled cavities [20], and so we shall neglect them here.

The c_{00} to which the $1S$ - $2S$ two-photon transition frequency is sensitive is the coefficient in the laboratory frame. In terms of the sun-centered coefficients, $c_{00} = c_{TT} + v_J c_{(TJ)}$, to first order in the velocity $\vec{v}$ of the laboratory. The c_{TT} contribution is uniform across all reference frames and therefore difficult to measure. However, it has been bounded at the 10^{-15} level using data from accelerator experiments [21, 22], which are more sensitive to violations of boost invariance. For these reasons, c_{TT} should be unimportant, and we

shall henceforth neglect it.

Since the contribution $c_{(TJ)}$ makes to c_{00} depends on the rest frame of the experiment, at different positions along the Earth's orbit the $c_{(TJ)}$ will affect atomic energy levels differently. Year-round measurements would make it possible to sample both $c_{(TJ)}$ corresponding to directions $\hat{J}$ lying in the ecliptic plane. Schematically, this means measuring a P -even observable in the laboratory frame, then comparing the measured values of this observable in laboratories moving with different velocities. The differences in measured values are not invariant under P and so are sensitive to the $c_{(TJ)}$, which are P -odd in the sun-centered frame.

The $c_{(TJ)}$ parameters describe violations of boost invariance. Testing this invariance requires a comparison of the physics in two different frames, and the effects of $c_{(TJ)}$ are suppressed by one power of the velocity difference between the frames being compared. In contrast, the isotropic boost invariance violation coefficient c_{TT} can only generate differences in physical observables between the two frames that are suppressed by two powers of the relative speed. It is probably because c_{TT} thus seems substantially more difficult to constrain on its own that the current methodology has not been much discussed. As well as mixing with c_{00} , the $c_{(TJ)}$ mix with the laboratory c_{jk} coefficients, which are relatively easy to bound using a rotating apparatus (or just the rotating Earth), and many attempts to constrain boost invariance violation have looked for differences in the anisotropy observed in different laboratory frames. If the pure anisotropy coefficients c_{JK} in the sun-centered frame are zero, then $c_{(jk)} = v_j c_{(Tk)} + v_k c_{(Tj)}$ in the laboratory frame. However, measurement of the anisotropy associated with this c_{jk} is not necessarily the best way to measure boost invariance violation. The mixings of the $c_{(TJ)}$ with both the c_{00} and c_{jk} generate observable effects at first order in $\vec{v}$.

The ^1H atom is an extremely clean system, containing only an electron and a proton. This simplifies theoretical calculations of energy eigenvalues, and it makes it easier to extract bounds on electron-sector coefficients; there are no many-body effects to complicate things. The presence of $c^{\nu\mu}$ shifts the energy of a nS state by

$$\Delta E = -\frac{m\alpha^2}{2n^2} \left(c_{00} + \frac{2}{3} c_{jj} \right); \quad (3)$$

as expected, only those laboratory-frame coefficients that are even under all discrete symmetries contribute. Moreover, because of the extremely long lifetime (≈ 0.122 s) of the ^1H $2S$ state, the $2S$ - $1S$ two-photon transition frequency can be measured quite precisely. These facts make this system an extremely attractive place to search for evidence of Lorentz violation.

Precisely the right kind of data for constraining the $c_{(TJ)}$ using ^1H spectrum measurements is already available. Recognizing the usefulness of the $2S$ - $1S$ ^1H transition as a probe of exotic physics, the laser spectroscopy group at the Max Planck Institute made multiple measurements of its frequency. They used these measurements to constrain the possible time variation of the fine structure constant α . However, their measurements,

separated by approximately 44 months, are also almost ideally suited to constrain the electron $c_{(TJ)}$, because the Earth's orbital velocities at the times of the two measurements were distinctly different.

The frequency difference between measurements made at two different points along the Earth's orbit is

$$\frac{\delta\nu}{\nu} = \frac{5}{3}\delta v_J c_{(TJ)}, \quad (4)$$

where $\delta\vec{v} = \vec{v}_2 - \vec{v}_1$ is the velocity difference between the two locations. Earth's velocity in its orbit is (neglecting small effects such as the orbit's ellipticity)

$$\vec{v} = v_{\oplus} \left(\hat{X} \sin \Omega_{\oplus} T - \hat{Y} \cos \eta \cos \Omega_{\oplus} T - \hat{Z} \sin \eta \cos \Omega_{\oplus} T \right), \quad (5)$$

where $v_{\oplus} \approx 10^{-4}$ and $\Omega_{\oplus}$ are the speed and angular frequency of the orbital revolution, and $\eta \approx 23.4^\circ$ is the inclination between the equatorial and ecliptic planes [23]. The additional velocity due to planetary rotation is substantially smaller.

The actual measurements were made over periods of weeks, during which time the velocity changed somewhat. In light of this fact, $\vec{v}_1$ and $\vec{v}_2$ should be replaced by average velocities over the periods during which the measurements were taken. If measurements are taken uniformly in time over a period ΔT , the average orbital velocity over this period is $\frac{\sin(\Omega_{\oplus}\Delta T/2)}{\Omega_{\oplus}\Delta T/2}$ times the velocity at the midpoint of the observation period. For a one-month observation window, the averaging reduces the velocity by less than 5%; for two months, the reduction is less than 18%. Averaging over June and July 1999 for the first measurement and February 2003 for the second, the effective velocity difference between the two observation periods is

$$\delta\vec{v} = -v_{\oplus}(1.35\hat{X} + 0.83\hat{Y} + 0.36\hat{Z}). \quad (6)$$

Measurements taken over short periods of time at opposite points on the orbit would maximize $|\delta\vec{v}|$, but the estimated experimental value (6) differs from the ideal one by only about 22%.

The observed difference $\delta\nu$ between the frequencies measured in the two experiments was (-29 ± 57) Hz. Compared with the $2S$ - $1S$ frequency of $2.4660611024748 \times 10^{15}$ Hz, this represents a fractional difference of $\frac{\delta\nu}{\nu} = (-1.2 \pm 2.4) \times 10^{-14}$. The resulting measurement of the electron $c_{(TJ)}$ coefficients is

$$0.83c_{(TX)} + 0.51c_{(TY)} + 0.22c_{(TZ)} = (4 \pm 8) \times 10^{-11}. \quad (7)$$

Another precision measurement of the $2S$ - $1S$ transition frequency ν would constrain an independent linear combination of the $c_{(TJ)}$ parameters. For optimal results, such a measurement should be performed in October or November, so the three measurements are roughly evenly spaced around the Earth's orbit. The linearly independent combination that would be bounded is $0.56c_{(TX)} - 0.76c_{(TY)} - 0.33c_{(TZ)}$. However, the third independent

combination of SME coefficients—which is the component of $c_{(TJ)}$ associated with boosts in the direction normal to the orbital plane—cannot be constrained this way, because the Earth’s velocity in this direction does not change. Laboratory measurements of this kind of boost invariance violation would need to rely on the Earth’s rotational velocity, which is about two orders of magnitude smaller than $\vec{v}_{\oplus}$ but which would combine with the Lorentz violation to produce sidereal variations in ν .

The $2S$ - $1S$ spectroscopy measurements used a ^{133}Cs atomic clock as a frequency standard. In principle, Lorentz violation in this nucleus could also affect the experimental results. However, the relevant nuclear SME parameters have already been strongly constrained [8] by a comparison of ^{133}Cs atomic clocks with other frequency standards. Ultimately, only differences between these frequency standards are observable, and the $2S$ - $1S$ spectroscopy provides a new constraint on the difference between the electron sector and others.

Other experiments already performed could provide similar but less precise bounds on the electron boost invariance violation coefficients. There are earlier, less precise measurements of the $2S$ - $1S$ frequency difference in ^1H [24]. There are also other recent measurements that were spread out in time to search for changes in α . Optical transitions in $^{199}\text{Hg}^+$ [25], $^{171}\text{Yb}^+$ [26], and ^{87}Sr [27] were studied over periods of years, with accuracies comparable to that of the ^1H experiment. These kinds of experiments could be used to place similar bounds, although the analyses are not so simple as with ^1H , since the transitions are no longer between pairs of isotropic states. The dependence on the laboratory c_{jk} coefficients would depend on the orientation of the apparatus, and the energies would be subject to sidereal variations. The atomic structures are also more complicated. Moreover, the $^{199}\text{Hg}^+$ and $^{171}\text{Yb}^+$ bounds were based on measurements made mostly around one time of year, which makes them less useful for constraining annual variations. The Sr experiment looked explicitly for annual variations, but the analysis assumed that frequency changes would be tied to variations in the Earth-sun distance. Taken together, the data from all these experiments suggest that both components of $c_{(TJ)}$ corresponding to directions $\hat{J}$ in the orbital plane should be bounded at the $\lesssim 10^{-9}$ level, but further analysis would be required to make this bound firm.

The measurement and remeasurement of the $2S$ - $1S$ two-photon transition frequency ν was conceived as a way to measure the time variation of α . But many of the same properties of these measurements—their high precision, relatively short duration, and substantial separation in time—that made them ideal for constraining $\dot{\alpha}$ also make them sensitive to the Lorentz violation coefficients $c_{(TJ)}$. The experimental data constrain a specific combination of these coefficients at the 10^{-11} level; this represents an eight order of magnitude improvement over previous laboratory constraints. Moreover, these bounds could easily be extended to another combination of $c_{(TJ)}$ coefficients simply by repeating the experimental measurement a third time.

References

- [1] For an overview, see *Proceedings of the Fourth Meeting on CPT and Lorentz Symmetry*, edited by V. A. Kostelecký (World Scientific, Singapore, 2008).
- [2] D. Colladay, V. A. Kostelecký, Phys. Rev. D **58**, 116002 (1998).
- [3] V. A. Kostelecký, N. Russell, arXiv:0801.0287.
- [4] C. J. Berglund, L. R. Hunter, D. Krause, Jr., E. O. Prigge, M. S. Ronfeldt, S. K. Lamoreaux, Phys. Rev. Lett. **75**, 1879 (1995).
- [5] V. A. Kostelecký, C. D. Lane, Phys. Rev. D **60**, 116010 (1999).
- [6] M. A. Humphrey, D. F. Phillips, E. M. Mattison, R. F. C. Vessot, R. E. Stoner, R. L. Walsworth, Phys. Rev. A **68**, 063807 (2003).
- [7] F. Canè, D. Bear, D. F. Phillips, M. S. Rosen, C. L. Smallwood, R. E. Stoner, R.L. Walsworth, V. A. Kostelecký, Phys. Rev. Lett. **93**, 230801 (2004).
- [8] P. Wolf, F. Chapelet, S. Bize, A. Clairon, Phys. Rev. Lett. **96**, 060801 (2006).
- [9] T. W. Kornack, G. Vasilakis, M. V. Romalis, in *Proceedings of the Fourth Meeting on CPT and Lorentz Symmetry*, edited by V. A. Kostelecký (World Scientific, Singapore, 2008).
- [10] M. Niering, *et al.*, Phys. Rev. Lett. **84**, 5496 (2000).
- [11] M. Fischer, *et al.*, Phys. Rev. Lett. **92**, 230802 (2004).
- [12] R. Bluhm, V. A. Kostelecký, N. Russell, Phys. Rev. Lett. **82**, 2254 (1999).
- [13] B. R. Heckel, C. E. Cramer, T. S. Cook, S. Schlamming, E. G. Adelberger, U. Schmidt, Phys. Rev. Lett. **97**, 021603 (2006).
- [14] V. A. Kostelecký and C. D. Lane, J. Math. Phys. **40**, 6245 (1999).
- [15] R. Bluhm, V. A. Kostelecký, C. D. Lane, N. Russell, Phys. Rev. D **68**, 125008 (2003).
- [16] C. D. Lane, Phys. Rev. D **72**, 016005 (2005).
- [17] G. Saathoff, S. Karpuk, U. Eisenbarth, G. Huber, S. Krohn, R. Muñoz Horta, S. Reinhardt, D. Schwalm, A. Wolf, G. Gwinner, Phys. Rev. Lett. **91**, 190403 (2003).
- [18] B. Altschul, Phys. Rev. D **74**, 083003 (2006).
- [19] B. Altschul, Astropart. Phys. **28**, 380 (2007).

- [20] H. Müller, Phys. Rev. D **71**, 045004 (2005).
- [21] M. A. Hohensee, R. Lehnert, D. F. Phillips, R. L. Walsworth, Phys. Rev. Lett. **102**, 170402 (2009).
- [22] B. Altschul, Phys. Rev D **80**, 091901(R) (2009).
- [23] V. A. Kostelecký, M. Mewes, Phys. Rev. D **66**, 056005 (2002).
- [24] Th. Udem, A. Huber, B. Gross, J. Reichert, M. Prevedelli, M. Weitz, and T. W. Hänsch, Phys. Rev. Lett. **79**, 2646 (1997).
- [25] S. Bize, *et al.*, Phys. Rev. Lett. **90**, 150802 (2003).
- [26] E. Peik, B. Lipphardt, H. Schnatz, T. Schneider, Chr. Tamm, S. G. Karshenboim, Phys. Rev. Lett. **93**, 170801 (2004).
- [27] S. Blatt, *et al.*, Phys. Rev. Lett. **100**, 140801 (2008).